

# PEDIATRICS®

## **Association Between Fetal Lymphedema and Congenital Cardiovascular Defects in Turner Syndrome**

Melissa L. Loscalzo, Phillip L. Van, Vincent B. Ho, Vladimir K. Bakalov, Douglas R.  
Rosing, Carol A. Malone, Harry C. Dietz and Carolyn A. Bondy

*Pediatrics* 2005;115;732-735

DOI: 10.1542/peds.2004-1369

**This information is current as of April 14, 2006**

The online version of this article, along with updated information and services, is  
located on the World Wide Web at:

<http://www.pediatrics.org/cgi/content/full/115/3/732>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2005 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# Association Between Fetal Lymphedema and Congenital Cardiovascular Defects in Turner Syndrome

Melissa L. Loscalzo, MD\*; Phillip L. Van, MS‡; Vincent B. Ho, MD§; Vladimir K. Bakalov, MD‡; Douglas R. Rosing, MD||; Carol A. Malone, PA-C||; Harry C. Dietz, MD\*¶; and Carolyn A. Bondy, MD‡

**ABSTRACT.** *Objectives.* Turner syndrome (TS) is associated with congenital cardiovascular defects (CCVDs), most commonly bicuspid aortic valve (BAV) and aortic coarctation (COARC), congenital renal anomalies, and fetal lymphedema. It has been theorized that compressive or obstructive effects of fetal lymphedema may actually cause cardiovascular and renal dysmorphogenesis in TS. The objective of this study was to determine whether there is a specific association between a history of fetal lymphedema and CCVDs in monosomy X, or TS, independent of karyotype or general severity of the phenotype.

*Methods.* This was a prospective study of 134 girls and women who have TS (mean age: 30 years) and were clinically evaluated for evidence of fetal lymphedema, classified as central (signified by the presence of neck webbing) or peripheral (current or perinatal, or dysplastic fingernails). The presence of BAV and/or COARC was detected by magnetic resonance imaging combined with echocardiography, and renal anomalies were determined by ultrasound.

*Results.* There is a strong association between developmental central lymphedema, signified by neck webbing, and the presence of BAV ( $\chi^2 = 10$ ) and COARC ( $\chi^2 = 8$ ). The association between webbed neck and CCVDs was independent of karyotype. There was, in contrast, no significant association between renal anomalies and webbed neck or CCVDs.

*Conclusions.* The strong, statistically significant association between neck webbing and the presence of BAV and COARC in TS suggests a pathogenetic connection between fetal lymphatic obstruction and defective aortic development. The presence of neck webbing in TS should alert the clinician to the possibility of congenital cardiovascular defects. *Pediatrics* 2005;115:732–735; *X-chromosome, aortic coarctation, bicuspid aortic valve, lymphedema.*

ABBREVIATIONS. CCVD, congenital cardiovascular defect; BAV,

From the \*McKusick-Nathans Institute of Genetic Medicine and †Howard Hughes Medical Institute, Johns Hopkins University School of Medicine, Baltimore, Maryland; ‡Developmental Endocrinology Branch, National Institute of Child Health, National Institutes of Health, Bethesda, Maryland; §Diagnostic Radiology Department, Warren G. Magnuson Clinical Center, National Institutes of Health, and Department of Radiology, Uniformed Services University of the Health Sciences, Bethesda, Maryland; ||National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, Maryland.

Dr Loscalzo and Mr Van contributed equally to this work.

Accepted for publication Jul 26, 2004.

doi:10.1542/peds.2004-1369

No conflict of interest declared.

Reprint requests to (C.A.B.) Building 10/10N262, National Institutes of Health, 10 Center Dr, Bethesda, MD 20892. E-mail: bondyc@mail.nih.gov  
PEDIATRICS (ISSN 0031 4005). Published in the public domain by the American Academy of Pediatrics.

bicuspid aortic valve; COARC, coarctation; TS, Turner syndrome.

Congenital cardiovascular defects (CCVDs), including most commonly bicuspid aortic valve (BAV) and coarctation of the aorta (COARC), are well-known features of Turner syndrome (TS).<sup>1–3</sup> It is unknown whether these defects are attributable to haploinsufficiency for X-chromosome gene(s) involved in cardiovascular development or secondary to other features of the syndrome, such as massive fetal lymphedema interfering with heart and major vessel formation. This latter possibility was suggested by observations of an apparent increased prevalence of CCVD in individuals with TS and neck webbing<sup>4</sup>—the postnatal residua of nuchal cystic hygromas caused by obstructed jugular lymphatics in utero. On the basis of this observation, Clark<sup>4</sup> proposed that jugular lymphatic obstruction led to compression of the ascending aorta, resulting in reduced flow and left-sided cardiovascular outflow tract anomalies. This view was supported by further epidemiologic observations in a study of 120 infants with neck webbing reported in the Iowa Birth Defects registry, among which 66% were found to have flow-related defects.<sup>5</sup> This hypothesis obtained additional support from studies of 45,X embryos ascertained because of cystic hygromas, with the majority demonstrating left-sided defects, from left heart hypoplasia to aortic valve defects, aortic hypoplasia, and/or coarctation.<sup>6,7</sup> Moreover, these studies documented the presence of dilated lymphatics in the vicinity of the developing heart and great vessels.

The original observations associating neck webbing with CCVDs came from a retrospective review of published cardiology clinic cases,<sup>4</sup> suggesting a likely bias toward the most severely affected individuals. Likewise, the pathologic studies certainly focused on the most severely affected fetuses, raising the possibility that the association between CCVDs and neck webbing simply reflects the most severe phenotype in 45,X individuals rather than a specific connection between these 2 phenotypic features of X-chromosome deletion. In the present study, we revisit the issue of association between signs of fetal lymphedema and the presence of CCVDs in a prospective study of 134 volunteers with TS, not selected for cardiovascular disease, who underwent physical examination and cardiovascular evaluation by MRI and echocardiography in a tertiary clinical research center.

## METHODS

Individuals with TS were recruited for this clinical study mainly through notices posted on Internet web sites (eg, <http://turners.nichd.nih.gov/>). Study participants signed informed assents/consents that were approved by the National Institute of Child Health and Human Development Institutional Review Board. Karyotype analysis of G-banded chromosomes in 50 lymphocytes was performed for all study participants. Diagnosis of TS was based on X-monosomy or X- or Y-deletion/rearrangement affecting  $\geq 70\%$  of cells. Participants with  $\geq 30\%$  normal cells were excluded from the study. A total of 134 girls and women with TS (mean age: 30; range: of 7–60 years) were studied. Adult participants underwent MRI that included contrast-enhanced, 3-dimensional MR angiography<sup>8</sup>; girls who were younger than 18 years had noncontrast MRI. All participants also had 2-D echocardiography with Doppler and renal ultrasound examination. MR images were evaluated by a single cardiovascular radiologist (V.B.H.), and echocardiograms were evaluated by a single cardiologist (D.R.R.). There was complete concordance for the 2 imaging methods for diagnosis of BAV, but echo was unable to define the aortic valve in 34 cases. MRI, however, was able to define the valve in all but 9 of these 34 patients, so the total for evaluation of BAV was 125. The diagnosis of COARC was by MRI or by history of surgical COARC repair.

Each study participant underwent a full physical examination and clinical cardiac evaluation. The presence of neck webbing was distinguished from a simply short neck by the finding of skin folds flaring from the lateral base of the skull to the mid- or lateral shoulder. The findings reported on physical examination were confirmed by review of neck photographs in anterior, lateral, and posterior views. A few participants had a history of surgical correction of webbing, verified by residual webbing and surgical scars.

Associations between phenotypic features were evaluated by  $\chi^2$  analysis with Yate's correction and forward stepwise regression in a linear model (SigmaStat, version 2.0; SPSS Inc, Chicago, IL).

## RESULTS

We found that 26 (21%) of 125 of our study participants had a BAV, and 16 (12%) of 134 had COARC, with 9 individuals having both. Forty-seven (35%) of 134 had a webbed neck. The association of webbed neck and these specific CCVDs was assessed by  $\chi^2$  analyses (Table 1). BAV was significantly associated with neck webbing, with a  $P < .002$ . COARC was also significantly associated with neck webbing, as was the presence of either COARC or BAV (Table 1). Of note, the presence of COARC was highly associated with BAV ( $\chi^2 = 15.8$ ,  $P = .0001$ ). Nine participants had both COARC and BAV, and 5 had COARC with a normal, tricuspid valve (1 had an aortic valve that was not defined by MR angiography or echocardiography).

It is often assumed that the more severe phenotypic features of TS may be associated simply because they are more frequently found in individuals with a 45,X karyotype. We found a strong statistical association between the presence of a webbed neck

and the 45,X karyotype ( $P < .001$ ) and a weaker but significant association between BAV and the 45,X karyotype ( $P = .014$ ). However, in a forward stepwise regression in a linear model with independent variables karyotype and webbed neck, only a webbed neck was a significant predictor of BAV ( $P < .0001$ ). Likewise, in the same type of regression analysis, webbed neck but not karyotype was significantly associated with COARC ( $P < .002$ ). We also investigated the relationship between CCVDs and peripheral lymphedema, which was identified in 2 ways: by a history of perinatal peripheral lymphedema and/or current peripheral lymphedema and by the presence of dysplastic fingernails. These diagnoses, each including 30% to 35% of study participants, were not associated with BAV, COARC, or webbed neck.

To clarify the specificity of the association between webbed neck and cardiovascular anomalies, we investigated the association between webbed neck and renal anomalies. Renal anomalies, including horseshoe kidney, single kidney, and duplicated collecting systems, were found in 28 (21%) of 134; 11 of these had a webbed neck, and 17 did not ( $\chi^2 = 0.385$ ,  $P = .65$ ). There was no association between the presence of BAV or COARC and renal anomalies (BAV:  $\chi^2 = 0.24$ ,  $P = .99$ ; COARC:  $\chi^2 = 1.8$ ,  $P = .2$ ). The presence of renal anomaly, however, was significantly associated with the 45,X karyotype ( $P < .02$ ).

The prevalence of congenital BAV or COARC by TS karyotype is shown in Table 2. This table is provided primarily as a reference for meta-analyses to accumulate data on genotype-phenotype correlations in larger numbers of subjects, because there are clearly too few individuals in genotype groups other than 45,X to allow any statistically meaningful inferences from this study alone. It is worth noting, however, that a relatively large percentage of 45,X/46,XY individuals have CCVDs. Adding our data to that previously reported, summarized by Sybert,<sup>3</sup> the prevalence of CCVDs in 45,X/46,XY is 14 (50%) of 28, substantially higher than the 30% to 40% typically noted in 45,X. To facilitate the localization of the CCVD locus on the X chromosome, we listed separately individuals with virtually complete deletions of Xp or Xq. We found BAV in 1 of 7 participants with 46,XiXq—which, combined with the 3 of 27 reported by Sybert for a prevalence of 12%, is significantly above that of the normal population, whereas to our knowledge, there are no reported congenital heart defects in subjects with 46,XXq-

**TABLE 1.** Association Between Webbed Neck and Cardiac Anomalies in TS

Webbed Neck	Aortic Valve			COARC+	COARC			BAV or COARC		
	BAV	TAV	Total		COARC-	Total	Either	Neither	Total	
Web+	16	28	44	11	36	47	18	26	44	
Web-	10	71	81	5	82	87	15	67	82	
Total	26	99	125	16	118	134	33	93	126	
$\chi^2$	10.0				9.05			8.5		
$P$	.002				.003			.004		
OR (95% CI)	4.1 (1.6–10.0)				5.0 (1.6–15.5)			3.3 (1.5–7.4)		

OR indicates odds ratio; CI, confidence interval.

**TABLE 2.** Genotype and Congenital Cardiovascular Defects (BAV or COARC)

	45X	45X/46XX*	45X/46XXp <sub>del</sub>	46X,Xp <sub>del</sub>	45X/46XY	46X,Xq <sub>del</sub>	45X/46XXq <sub>del</sub>	Total
N	83	21	14	8	5	2	2	134
No with cardiac defects	27	0	2	1	3	0	1	34
Incidence of cardiac defects, %	33	0	14	13	60	0	50	25

X\* indicates a normal X, ring X, or X with partial q or p deletions.  
Xp<sub>del</sub> includes only the most proximal small arm deletions (<11.3) and iXq.

## DISCUSSION

The present observations in a large group of participants with TS not selected for cardiovascular disease confirms a significant association between central fetal lymphedema, signaled by neck webbing, BAV, and COARC. We have also shown that the association between neck webbing and CCVDs is not merely a reflection of a more severe phenotype in 45,X individuals but is a statistically significant connection independent of karyotype and parental imprinting and specific for cardiovascular but not renal anomalies. Our study confirms the modern observation of Mazzanti et al<sup>2</sup> that COARC is significantly associated with neck webbing and extends the finding to BAV as well. This highly significant association between central lymphedema during fetal development and CCVDs could reflect a cause and effect relation between these 2 features.

One study suggested that defective cardiovascular development may be a primary feature in TS, with impaired lymphatic development and generalized hydrops as consequences.<sup>9</sup> This proposition seems unlikely for a number of reasons. The anatomic defects associated with fetal lymphedema in TS are decreased numbers of lymphatics and dilated lymphatic channels that end in distended sacs, lacking connections with the venous system.<sup>4,6,10,11</sup> The developmental delay is particularly pronounced in the axillojugular lymphatics,<sup>10</sup> which drain the upper half of the body. The nuchal cystic hygromas are loculated collections of lymph associated with blind-ended jugular lymphatics. Central lymphatic obstruction is apparent in fetuses with TS as early as 10 to 12 weeks but resolves during the latter half of gestation and is detected at birth only by the redundant skin folds of the neck. It is difficult to imagine how delayed or defective lymphatic development could be caused by heart failure as shown in the Barr and Oman-Ganes study.<sup>9</sup> It is equally difficult to reconcile this notion with the observation of many individuals who have TS with neck webbing, with or without CCVDs, but no sign of heart failure. It seems more likely that severe lymphatic obstruction early in fetal development may cause heart failure from compression and/or impaired filling of developing cardiovascular structures, leading to fetal hydrops and demise, as was the case for most of the cases examined in the Barr and Oman-Ganes study.<sup>9</sup> Thus, if there were a cause-and-effect relation between fetal lymphedema and CCVDs in TS, then it would seem most likely that the lymphedema is primary.

Clarke<sup>4</sup> suggested that centrally localized distended lymphatics compress the developing aortic root, resulting in specific left-sided defects, including hypoplastic left heart, BAV, and COARC as a result

of low flow, and specific right-sided defects such as persistent left superior vena cava, anomalous pulmonary venous return, and dilated right atrium as a result of back-pressure from obstruction to forward flow. The Clarke hypothesis seems plausible and does include the constellation of CCVDs most commonly found in TS, but after 20 years, it remains hypothetical, without direct proof. Although anatomic evidence shows that dilated lymphatics are observed in the vicinity of developing aorta, no correlation has been observed between the presence of dilated lymphatic and aortic defects.<sup>4,6,7</sup> The present study provides a significant correlation between the presence of CCVDs and centralized fetal lymphedema but obviously cannot establish cause and effect.

An alternative explanation for the association between neck webbing and CCVDs in TS is that haploinsufficiency for an X-chromosome gene independently causes fetal lymphedema (webbed neck) and congenital heart defects. Haploinsufficiency for an autosomal gene (*FOXC2*;16q) causes widespread lymphedema and occasional cardiac defects in the lymphedema-distichiasis syndrome,<sup>12</sup> but targeted deletion of this gene in mice results in abnormal aortic arch development in the absence of obvious lymphedema.<sup>13</sup> These observations suggest that the heart defects and lymphedema found in the human disorder may be independent effects of the same gene. Noonan syndrome may be another instance in which lymphedema and cardiac defects occur as independent effects of the same gene mutation/deletion. Heart defects are more common and include predominantly right-sided lesions in Noonan syndrome compared with TS.<sup>14</sup> Also in contrast to TS, Noonan syndrome manifests few abnormal features during early fetal development, with cystic hygroma being a rare finding and residual stigmata of fetal lymphedema are less common in Noonan syndrome.<sup>15-17</sup> Attempts to correlate webbed neck and cardiovascular lesions in this syndrome have been conflicting.<sup>5,18</sup> A specific gene mutation that is responsible for ~50% of Noonan syndrome cases has only recently been identified<sup>19</sup>; thus, the earlier studies that attempted to associate CCVDs and lymphedema may have included genetically heterogeneous populations.

The role of fetal lymphedema in the pathogenesis of CCVDs in TS may be resolved when the genes that are responsible for lymphedema are identified, allowing experimental studies to elucidate its function in developing lymphatic and cardiovascular systems. With this goal, a recent study has suggested that the "critical region" for the TS lymphedema gene lies at Xp11.4.<sup>20</sup> The careful phenotypic characterization of additional patients with TS and infor-

mative X-chromosome deletions will further advance this effort, aided by newly available X-chromosome sequence information.

### ACKNOWLEDGMENTS

We are grateful to Eileen Lange, RN, and the nurses of the National Institutes of Health Clinical Center for excellent care of our patients. We also acknowledge the important role of the Turner Syndrome Society, USA, in publicizing the study.

### REFERENCES

1. Gotzsche CO, Krag-Olsen B, Nielsen J, Sorensen KE, Kristensen BO. Prevalence of cardiovascular malformations and association with karyotypes in Turner's syndrome. *Arch Dis Child.* 1994;71:433-436
2. Mazzanti L, Cacciari E. Congenital heart disease in patients with Turner's syndrome. Italian Study Group for Turner Syndrome (ISGTS). *J Pediatr.* 1998;133:688-692
3. Sybert VP. Cardiovascular malformations and complications in Turner syndrome. *Pediatrics* 1998;101(1). Available at: [www.pediatrics.org/cgi/content/full/101/1/e11](http://www.pediatrics.org/cgi/content/full/101/1/e11)
4. Clark EB. Neck web and congenital heart defects: a pathogenic association in 45 X-O Turner syndrome. *Teratology.* 1984;29:355-361
5. Berdahl LD, Wenstrom KD, Hanson JW. Web neck anomaly and its association with congenital heart disease. *Am J Med Genet.* 1985;56:304-307
6. Lacro RV, Jones KL, Benirschke K. Coarctation of the aorta in Turner syndrome: a pathologic study of fetuses with nuchal cystic hygromas, hydrops fetalis, and female genitalia. *Pediatrics.* 1988;81:445-451
7. Miyabara S, Nakayama M, Suzumori K, Yonemitsu N, Sugihara H. Developmental analysis of cardiovascular system of 45,X fetuses with cystic hygroma. *Am J Med Genet.* 1997;68:135-141
8. Ho V, Prince M. Thoracic MR aortography: imaging techniques and strategies. *Radiographics.* 1998;18:287-309
9. Barr M Jr, Oman-Ganes L. Turner syndrome morphology and morphometrics: cardiac hypoplasia as a cause of midgestation death. *Teratology.* 2002;66:65-72
10. van der Putte SC. Lymphatic malformation in human fetuses. A study of fetuses with Turner's syndrome or status Bonnevie-Ullrich. *Virchows Arch A Pathol Anat Histol.* 1977;376:233-246
11. Miyabara S, Sugihara H, Maehara N, et al. Significance of cardiovascular malformations in cystic hygroma: a new interpretation of the pathogenesis. *Am J Med Genet.* 1989;34:489-501
12. Fang J, Dagenais SL, Erickson RP, et al. Mutations in FOXC2 (MFH-1), a forkhead family transcription factor, are responsible for the hereditary lymphedema-distichiasis syndrome. *Am J Med Genet.* 2000;67:1382-1388
13. Iida K, Koseki H, Kakinuma H, et al. Essential roles of the winged helix transcription factor MFH-1 in aortic arch patterning and skeletogenesis. *Development.* 1997;124:4627-4638
14. Sarkozy A, Conti E, Seripa D, et al. Correlation between PTPN11 gene mutations and congenital heart defects in Noonan and LEOPARD syndromes. *J Med Genet.* 2003;40:704-708
15. Nisbet D, Griffin D, Chitty L. Prenatal features of Noonan syndrome. *Prenat Diagn.* 1999;19:642-647
16. Achiron R, Heggesh J, Grisaru D, et al. Noonan syndrome: a cryptic condition in early gestation. *Am J Med Genet.* 2000;92:159-165
17. Menashe M, Arbel R, Raveh D, Achiron R, Yagel S. Poor prenatal detection rate of cardiac anomalies in Noonan syndrome. *Ultrasound Obstet Gynecol.* 2002;19:51-55
18. Brady A, Patton M. Web-neck anomaly and its association with congenital heart disease. *Am J Med Genet.* 1996;64:605
19. Tartaglia M, Mehler E, Goldberg R, et al. Mutations in PTPN11, encoding the protein tyrosine phosphatase SHP-2, cause Noonan syndrome. *Nat Genet.* 2001;29:465-468
20. Boucher CA, Sargent CA, Ogata T, Affara NA. Breakpoint analysis of Turner patients with partial Xp deletions: implications for the lymphoedema gene location. *J Med Genet.* 2001;38:591-598

## THE COST OF MEDICAL EDUCATION

"The cost of obtaining a medical education has been spiraling upward for the past 20 years. . . . The average tuition and fees at public medical schools during the 2003-2004 academic year amounted to \$16 153, and the corresponding figure for private schools was \$32 588. Adding \$20 000 to \$25 000 for living expenses, books, and equipment brings the estimated cost of 4 years of attendance to about \$140 000 for public schools and \$225 000 for private schools. . . . The average debt carried by 1984 graduates was \$22 000 for public school and \$26 500 for private school. By 2004, the debt had increased to \$105 000 for public school and \$140 000 for private school, and only about 20% of medical students graduated with no debt."

Morrison G. Mortgaging our future: the cost of medical education. *N Engl J Med.* 2005;352:177-179

Noted by JFL, MD

## Association Between Fetal Lymphedema and Congenital Cardiovascular Defects in Turner Syndrome

Melissa L. Loscalzo, Phillip L. Van, Vincent B. Ho, Vladimir K. Bakalov, Douglas R. Rosing, Carol A. Malone, Harry C. Dietz and Carolyn A. Bondy

*Pediatrics* 2005;115:732-735

DOI: 10.1542/peds.2004-1369

**This information is current as of April 14, 2006**

<b>Updated Information &amp; Services</b>	including high-resolution figures, can be found at: <a href="http://www.pediatrics.org/cgi/content/full/115/3/732">http://www.pediatrics.org/cgi/content/full/115/3/732</a>
<b>References</b>	This article cites 19 articles, 6 of which you can access for free at: <a href="http://www.pediatrics.org/cgi/content/full/115/3/732#BIBL">http://www.pediatrics.org/cgi/content/full/115/3/732#BIBL</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Heart &amp; Blood Vessels</b> <a href="http://www.pediatrics.org/cgi/collection/heart_and_blood_vessels">http://www.pediatrics.org/cgi/collection/heart_and_blood_vessels</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.pediatrics.org/misc/Permissions.shtml">http://www.pediatrics.org/misc/Permissions.shtml</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://www.pediatrics.org/misc/reprints.shtml">http://www.pediatrics.org/misc/reprints.shtml</a>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

